Brain Research, 507 (1990) 357–359
Elsevier

Author Index

Agid, Y., see Raisman-Vozari, R., 45
Akaike, N., see Maruyama, T., 17
Akimoto, K., Hamamura, T., Kazahaya, Y., Akiyama, K. and Otsuki, S., Enhanced extracellular dopamine level may be the fundamental neuropharmacological basis of cross-behavioral sensitization between methamphetamine and cocaine — an in vivo dialysis study in freely moving rats, 344
Akiyama, H., see Kaneko, T., 151
Akiyama, K., see Akimoto, K., 344
Assadian, H., Koshiyama, H., 321
Atchison, N.E., see Osgood, P.F., 11
Azmitia, E.C., see Zhou, F.C., 301

Bak, T.H., see Carr, K.D., 289
Bakshi, R. and Faden, A.I., Competitive and non-competitive NMDA antagonists limit dynorphin A-induced rat hindlimb paralysis, 1
Barnes, J.M., Barnes, N.M., Costall, B., Horovitz, Z.P., Ironside, J.W., Naylor, R.J. and Williams, T.J., Angiotensin II inhibits acetylcholine release from human temporal cortex: implications for cognition, 341
Barnes, N.M., see Barnes, J.M., 341
Barraco, R.A., Janusz, C.A., Schoener, E.P. and Simpson, L.L., Cardiorespiratory function is altered by picomole injections of 5′-*N*-ethylcarboxamidoadenosine into the nucleus tractus solitarius of rats, 234
Berg-Johnsen, J. and Langmoen, I.A., Mechanisms concerned in the direct effect of isoflurane on rat hippocampal and human neocortical neurons, 28
Bianchi, A.L., see Iscoe, S., 281
Biegon, A., see Gross-Isseroff, R., 208
Blair-West, J.R., Denton, D.A., Gibson, A.P. and McKinley, M.J., Opening the blood–brain barrier, 6
Blendy, J.A., see Wahlestedt, C., 65
Bohan, M.C., see Gintzler, A.R., 312

Carr, D.B., see Osgood, P.F., 11
Carr, K.D. and Bak, T.H., Rostral and caudal ventricular infusion of antibodies to dynorphin A(1–17) and dynorphin A(1–8): effects on electrically-elicited feeding in the rat, 289
Costall, B., see Barnes, J.M., 341
Crockett, D.P., see Maslany, S., 164

Davis, K.L., see Haroutunian, V., 261
Delree, P., see Lefebvre, P.P., 254
Denton, D.A., see Blair-West, J.R., 6
DeTeresa, R., see Iimoto, D.S., 273
DeToledo-Morrell, L., see Geinisman, Y., 325
Dragunow, M., see Jansen, K.L.R., 158

Egger, M.D., see Maslany, S., 164
Ekman, R., see Wahlestedt, C., 65
Etgen, A.M., see Fleischmann, A., 161

Faden, A.I., see Bakshi, R., 1
Fastbom, J. and Fredholm, B.B., Effects of long-term theophylline treatment on adenosine A_1-receptors in rat brain: autoradiographic evidence for increased receptor number and altered coupling to G-proteins, 195
Faull, R.L.M., see Jansen, K.L.R., 158
Feder, H.H., see McCarthy, M.M., 40
Feuerstein, C., see Raisman-Vozari, R., 45
Fleischmann, A., Makman, M.H. and Etgen, A.M., Ovarian steroids increase veratridine-induced release of amino acid neurotransmitters in preoptic area synaptosomes, 161
Fredholm, B.B., see Fastbom, J., 195
Fujise, N., see Nakahara, N., 172

Geinisman, Y., Morrell, F. and DeToledo-Morrell, L., Increase in the relative proportion of perforated axospinous synapses following hippocampal kindling is specific for the synaptic field of stimulated axons, 325
Gibson, A.P., see Blair-West, J.R., 6
Gintzler, A.R. and Bohan, M.C., Pain thresholds are elevated during pseudopregnancy, 312
Girault, J.-A., see Raisman-Vozari, R., 45
Gotoh, M., see Iguchi, A., 295
Grélot, L., see Iscoe, S., 281
Grimelius, L., see Nässel, D.R., 225
Gross-Isseroff, R., Salama, D., Israeli, M. and Biegon, A., Autoradiographic analysis of [^{3}H]ketanserin binding in the human brain postmortem: effect of suicide, 208

Halsey, M.J., see Millan, M.H., 354
Hamamura, T., see Akimoto, K., 344
Haroutunian, V., Kanof, P.D., Tsuboyama, G. and Davis, K.L., Restoration of cholinomimetic activity by clonidine in cholinergic plus noradrenergic lesioned rats, 261
Harris, V., see Vasko, M.R., 129
Hartwieg, E.A., see Potrebic, S., 317
Hattori, N., see Koshiyama, H., 321
Heilig, M., see Wahlestedt, C., 65
Henry, J.L., see Salter, M.W., 69
Hoffman, P.L., see Ishizawa, H., 189
Hong, J.S., see Jiang, H.-K., 57
Höög, A., see Nässel, D.R., 225
Horovitz, Z.P., see Barnes, J.M., 341
Hultborn, H., see Kernell, D., 176

Ichikawa, J. and Meltzer, H.Y., Apomorphine does not reverse reduced basal dopamine release in rat striatum and nucleus accumbens after chronic haloperidol treatment, 138
Igisu, H., see Matsuoka, M., 351
Iguchi, A., Yatomi, A., Gotoh, M., Matsunaga, H., Uemura, K., Miura, H., Satake, T., Tamagawa, T. and Sakamoto, N., Neostigmine-induced hyperglycemia is mediated by central muscarinic receptor in fed rats, 295
Iimoto, D.S., Masliah, E., DeTeresa, R., Terry, R.D. and Saitoh, T., Aberrant casein kinase II in Alzheimer's disease, 273
Ikemoto, Y., see Maruyama, T., 17
Imai, S., see Onozuka, M., 143
Imura, H., see Koshiyama, H., 321
Inoue, M., see Kaneto, H., 35
Inoue, N., see Matsuoka, M., 351
Ironside, J.W., see Barnes, J.M., 341
Iscoe, S., Grélot, L. and Bianchi, A.L., Responses of inspiratory neurons of the dorsal respiratory group to stimulation of expiratory muscle and vagal afferents, 281
Ishikawa, Y., see Koshiyama, H., 321
Ishizawa, H., Tabakoff, B., Mefford, I.N. and Hoffman, P.L., Reduction of arginine vasopressin binding sites in mouse lateral septum by treatment with 6-hydroxydopamine, 189
Israeli, M., see Gross-Isseroff, R., 208
Iwata, K., Muramatsu, H., Tsuboi, Y. and Sumino, R., Input-output relationships in the jaw and orofacial motor zones of the cat cerebral cortex, 337

Jansen, K.L.R., Dragunow, M. and Faull, R.L.M., Sigma receptors are highly concentrated in the rat pineal gland, 158
Janusz, C.A., see Barraco, R.A., 234
Jenner, P., see Raisman-Vozari, R., 45
Jiang, H.-K., McGinty, J.F. and Hong, J.S., Differential modulation of striatonigral dynorphin and enkephalin by dopamine receptor subtypes, 57

Kamsteeg, H., see Wolterink, G., 92
Kamsteeg, H., see Wolterink, G., 101
Kaneko, T., Akiyama, H., Nagatsu, I. and Mizuno, N., Immunohistochemical demonstration of glutaminase in catecholaminergic and serotonergic neurons of rat brain, 151
Kaneto, H. and Inoue, M., Action site of adrenergic blockers to suppress the development of tolerance to morphine analgesia, 35
Kanof, P.D., see Haroutunian, V., 261
Kato, Y., see Koshiyama, H., 321
Kawanishi, G., see Nakahara, N., 172
Kazahaya, Y., see Akimoto, K., 344
Kazianis, A., see Osgood, P.F., 11
Kellar, K.J., see Wahlestedt, C., 65
Kemp, J.W., see Osgood, P.F., 11
Kernell, D. and Hultborn, H., Synaptic effects on recruitment gain: a mechanism of importance for the input-output relations of motoneurone pools?, 176
Kimura, I., Kondoh, T. and Kimura, M.,

Postsynaptic nicotinic ACh receptor-operated Ca^{2+} transients with neostigmine in phrenic nerve-diaphragm muscles of mice, 309
Kimura, M., see Kimura, I., 309
Kondoh, T., see Kimura, I., 309
Koshiyama, H., Shimatsu, A., Kato, Y., Assadian, H., Hattori, N., Ishikawa, Y., Tanoh, T., Yanaihara, N. and Imura H., Galanin-induced prolactin release in rats: pharmacological evidence for the involvement of α-adrenergic and opioidergic mechanisms, 321

Langmoen, I.A., see Berg-Johnsen, J., 28
Lechan, R.M., see Schipper, H.M., 200
Lefebvre, P.P., Leprince, P., Weber, T., Rigo, J.-M., Delree, P. and Moonen, G., Neuronotrophic effect of developing otic vesicle on cochleo-vestibular neurons: evidence for nerve growth factor involvement, 254
Leprince, P., see Lefebvre, P.P., 254
Leventhal, T.O., see Schwartzbaum, J.S., 85
Liebeskind, J.C., see Marek, P., 155
Lin, J., see Matsuoka, M., 351
Lovinger, D.M., see White, G., 332
Lundquist, T., see Nässel, D.R., 225

Maciewicz, R., see Potrebic, S., 317
Maeda, M., see Nakai, M., 168
Makman, M.H., see Fleischmann, A., 161
Malik, K.F., see McCarthy, M.M., 40
Marek, P., Yirmiya, R. and Liebeskind, J.C., Genetic influences on brain stimulation-produced analgesia in mice: II. Correlation with brain opiate receptor concentration, 155
Marsden, C.D., see Raisman-Vozari, R., 45
Maruyama, T., Ikemoto, Y. and Akaike, N., Effect of temperature on the inhibition of the GABA-gated response by intracellular calcium, 17
Maslany, S., Crockett, D.P. and Egger, M.D., Somatotopic organization of the cuneate nucleus in the rat: transganglionic labelling with WGA-HRP, 164
Masliah, E., see Iimoto, D.S., 273
Matsunaga, H., see Iguchi, A., 295
Matsuoka, M., Igisu, H., Lin, J. and Inoue, N., Effects of acrylamide and N,N′-methylene-bis-acrylamide on creatine kinase activity, 351
McCarthy, M.M., Malik, K.F. and Feder, H.H., Increased GABAergic transmission in medial hypothalamus facilitates lordosis but has the opposite effect in preoptic area, 40
McGinty, J.F., see Jiang, H.-K., 57
McKinley, M.J., see Blair-West, J.R., 6
Mefford, I.N., see Ishizawa, H., 189
Meldrum, B.S., see Millan, M.H., 354
Meltzer, H.Y., see Ichikawa, J., 138
Millan, M.H., Wardley-Smith, B., Halsey, M.J. and Meldrum, B.S., Effect of NMDA and 2-amino-7-phosphonoheptanoate focal injection into the ventrolateral thalamic nucleus on the high pressure neurological syndrome in the rat, 354
Miura, H., see Iguchi, A., 295
Miura, M., see Saji, M., 247
Mizobe, F., see Nakahara, N., 172
Mizuno, N., see Kaneko, T., 151
Moonen, G., see Lefebvre, P.P., 254
Morrell, F., see Geinisman, Y., 325
Moussaoui, S., see Raisman-Vozari, R., 45
Muramatsu, H., see Iwata, K., 337

Nadakavukaren, J.J., Welsh, D.K. and Reppert, S.M., Aluminum fluoride reveals a phosphoinositide system within the suprachiasmatic region of rat hypothalamus, 181
Nagatsu, I., see Kaneko, T., 151
Nakahara, N., Fujise, N., Kawanishi, G. and Mizobe, F., Central muscarinic activities of an M_1-selective agonist: preferential effect on reversal of amnesia, 172
Nakai, M., Tamaki, K., Yamamoto, J., Shimouchi, A. and Maeda, M., A minimally invasive technique for multiple measurement of regional blood flow of the rat brain using radiolabeled microspheres, 168
Nässel, D.R., Lundquist, T., Höög, A. and Grimelius, L., Substance P-like immunoreactive neurons in the nervous system of *Drosophila,* 225
Naylor, R.J., see Barnes, J.M., 341
Nishimura, S., see Sorimachi, M., 347

O'Neill, R.D., Uric acid levels and dopamine transmission in rat striatum: diurnal changes and effects of drugs, 267
Onozuka, M. and Imai, S., Induction of epileptic seizure activity by a specific protein from cobalt-induced epileptogenic cortex of rats, 143
Osgood, P.F., Carr, D.B., Kazianis, A., Kemp, J.W., Atchison, N.E. and Szyfelbein, S.K., Antinociception in the rat induced by a cold environment, 11
Otsuki, S., see Akimoto, K., 344

Plenderleith, M.B. and Snow, P.J., The effect of peripheral nerve section on lectin binding in the superficial dorsal horn of the rat spinal cord, 146
Potrebic, S., Strassman, A., Hartwieg, E.A. and Maciewicz, R., Ultrastructure of intracellularly labeled trigeminal vascular convergence neurons, 317

Radhakishun, F.S., see Wolterink, G., 92
Radhakishun, F.S., see Wolterink, G., 101
Raisman-Vozari, R., Girault, J.-A., Moussaoui, S., Feuerstein, C., Jenner, P., Marsden, C.D. and Agid, Y., Lack of change in striatal DARPP-32 levels following nigrostriatal dopaminergic lesions in animals and in parkinsonian syndromes in man, 45
Reichlin, S., see Schipper, H.M., 200
Reppert, S.M., see Nadakavukaren, J.J., 181
Rigo, J.-M., see Lefebvre, P.P., 254

Saitoh, T., see Iimoto, D.S., 273
Saji, M. and Miura, M., Thoracic expiratory motor neurons of the rat: localization and sites of origin of their premotor neurons, 247
Sakamoto, N., see Iguchi, A., 295
Salama, D., see Gross-Isseroff, R., 208
Salter, M.W. and Henry, J.L., Physiological characteristics of responses of wide dynamic range spinal neurones to cutaneously applied vibration in the cat, 69
Satake, T., see Iguchi, A., 295
Schipper, H.M., Lechan, R.M. and Reichlin, S., Glial peroxidase activity in the hypothalamic arcuate nucleus: effects of estradiol valerate-induced persistent estrus, 200
Schoener, E.P., see Barraco, R.A., 234
Scholfield, C.N., Properties of K-currents in unmyelinated presynaptic axons of brain revealed by extracellular polarisation, 121
Schwartzbaum, J.S. and Leventhal, T.O., Neural substrates of behavioral aversion in lateral hypothalamus of rabbits, 85
Shaikh, M.B. and Siegel, A., GABA-mediated regulation of feline aggression elicited from midbrain periaqueductal gray, 51
Shimatsu, A., see Koshiyama, H., 321
Shimouchi, A., see Nakai, M., 168
Siegel, A., see Shaikh, M.B., 51
Simpson, L.L., see Barraco, R.A., 234
Snow, P.J., see Plenderleith, M.B., 146
Sorimachi, M., Yamagami, K. and Nishimura, S., Tetraethylammonium stimulates adrenomedullary secretion by causing fluctuations in a cytosolic free Ca concentration, 347
Speth, R.C., see Walters, D.E., 23
Strassman, A., see Potrebic, S., 317
Sumino, R., see Iwata, K., 337
Szyfelbein, S.K., see Osgood, P.F., 11

Tabakoff, B., see Ishizawa, H., 189
Tamagawa, T., see Iguchi, A., 295
Tamaki, K., see Nakai, M., 168
Tanoh, T., see Koshiyama, H., 321
Terry, R.D., see Iimoto, D.S., 273
Tsuboi, Y., see Iwata, K., 337
Tsuboyama, G., see Haroutunian, V., 261

Uemura, K., see Iguchi, A., 295
Umetani, T., Topographic organization of the cerebellar nucleocortical projection in the albino rat: an autoradiographic orthograde study, 216

Van Ree,, J.M., see Wolterink, G., 92
Van Ree, J.M., see Wolterink, G., 101
Van Ree, J.M., see Wolterink, G., 109
Van Ree, J.M., see Wolterink, G., 115
Van Zanten, E., see Wolterink, G., 92
Van Zanten, E., see Wolterink, G., 101
Van Zanten, E., see Wolterink, G., 115
Vasko, M.R. and Harris V., γ-Aminobutyric acid inhibits the potassium-stimulated release of somatostatin from rat spinal cord slices, 129

Wahlestedt, C., Blendy, J.A., Kellar, K.J., Heilig, M., Widerlöv, E. and Ekman, R., Electroconvulsive shocks increase the concentration of neocortical hippocampal neuropeptide Y (NPY)-like immunoreac-

tivityin the rat, 65
Walters, D.E. and Speth, R.C., Loss of striatal angiotensin-converting enzyme following intrastriatal AF64A is not related to destruction of cholinergic interneurons, 23
Wardley-Smith, B., see Millan, M.H., 354
Weber, T., see Lefebvre, P.P., 254
Weight, F.F., see White, G., 332
Welsh, D.K., see Nadakavukaren, J.J., 181
White, G., Lovinger, D.M. and Weight, F.F., Ethanol inhibits NMDA-activated current but does not alter GABA-activated current in an isolated adult mammalian neuron, 332
Widerlöv, E., see Wahlestedt, C., 65
Williams, T.J., see Barnes, J.M., 341
Wolterink, G., Van Zanten, E., Kamsteeg, H., Radhakishun, F.S. and Van Ree, J.M., Functional recovery after destruction of dopamine systems in the nucleus accumbens of rats. I. Behavioral and biochemical studies, 92
Wolterink, G., Van Zanten, E., Kamsteeg, H., Radhakishun, F.S. and Van Ree, J.M., Functional recovery after destruction of dopamine systems in the nucleus accumbens of rats. II. Facilitation by the ACTH-(4–9) analog ORG 2766, 101
Wolterink, G. and Van Ree, J.M., Functional recovery after destruction of dopamine systems in the nucleus accumbens of rats. III. Further analysis of the facilitating effect of the ACTH-(4–9) analog ORG 2766, 109
Wolterink, G., Van Zanten, E. and Van Ree, J.M., Functional recovery after destruction of dopamine systems in the nucleus accumbens of rats. IV. Delay by intra-accumbal treatment with ORG 2766- or α-MSH antiserum, 115

Yamagami, K., see Sorimachi, M., 347
Yamamoto, J., see Nakai, M., 168
Yanaihara, N., see Koshiyama, H., 321
Yatomi, A., see Iguchi, A., 295
Yirmiya, R., see Marek, P., 155

Zhou, F.C. and Azmitia, E.C., Neurotrophic factor for serotonergic neurons prevents degeneration of grafted raphe neurons in the cerebellum, 301